


- 38 -

CLAIMS:

1. A bleaching composition comprising:

a) a monomer ligand or transition metal catalyst thereof of
 5 a ligand having the formula (I):



wherein each R is independently selected from: hydrogen, F,
 10 Cl, Br, hydroxyl, C1-C4-alkylo-, -NH-CO-H, -NH-CO-C1-C4-
 alkyl, -NH₂, -NH-C1-C4-alkyl, and C1-C4-alkyl;

R1 and R2 are independently selected from:

C1-C4-alkyl,

C6-C10-aryl, and,

15 a group containing a heteroatom capable of coordinating to a
 transition metal, wherein at least one of R1 and R2 is the
 group containing the heteroatom;

R3 and R4 are independently selected from hydrogen, C1-C8
 alkyl, C1-C8-alkyl-O-C1-C8-alkyl, C1-C8-alkyl-O-C6-C10-aryl,
 20 C6-C10-aryl, C1-C8-hydroxyalkyl, and -(CH₂)_nC(O)OR₅

wherein R₅ is independently selected from: hydrogen, C1-C4-
 alkyl, n is from 0 to 4, and mixtures thereof; and,

X is selected from C=O, -[C(R₆)₂]_y- wherein Y is from 0 to 3
 each R₆ is independently selected from hydrogen, hydroxyl,

25 C1-C4-alkoxy and C1-C4-alkyl; and,

- 39 -

b) the balance carriers and adjunct ingredients.

2. A bleaching composition according to claim 1, wherein R1 and R2 are both selected from a group containing a

5 heteroatom capable of coordinating to a transition metal.

3. A bleaching composition according to claim 1, wherein the group containing the heteroatom is:

a heterocycloalkyl: selected from the group consisting of:

10 pyrrolinyl; pyrrolidinyl; morpholinyl; piperidinyl;
piperazinyl; hexamethylene imine; 1,4-piperazinyl;
tetrahydrothiophenyl; tetrahydrofuranlyl; tetrahydropyranyl;
and oxazolidinyl, wherein the heterocycloalkyl may be
connected to the ligand via any atom in the ring of the
15 selected heterocycloalkyl,

a -C1-C6-alkyl-heterocycloalkyl, wherein the
heterocycloalkyl of the -C1-C6-heterocycloalkyl is selected
from the group consisting of: piperidinyl; piperidine; 1,4-
piperazine, tetrahydrothiophene; tetrahydrofuran;

20 pyrrolidine; and tetrahydropyran, wherein the
heterocycloalkyl may be connected to the -C1-C6-alkyl via
any atom in the ring of the selected heterocycloalkyl,
a -C1-C6-alkyl-heteroaryl, wherein the heteroaryl of the -
C1-C6-alkylheteroaryl is selected from the group consisting

25 of: pyridinyl; pyrimidinyl; pyrazinyl; triazolyl;
pyridazinyl; 1,3,5-triazinyl; quinolinyl; isoquinolinyl;
quinoxalinyl; imidazolyl; pyrazolyl; benzimidazolyl;
thiazolyl; oxazolidinyl; pyrrolyl; carbazolyl; indolyl; and
isoindolyl, wherein the heteroaryl may be connected to the -
30 C1-C6-alkyl via any atom in the ring of the selected

- 40 -

heteroaryl and the selected heteroaryl is optionally substituted by -C1-C4-alkyl,

a -C0-C6-alkyl-phenol or thiophenol,

a -C2-C4-alkyl-thiol, thioether or alcohol,

5 a -C2-C4-alkyl-amine, and

a -C2-C4-alkyl-carboxylate.

4. A bleaching composition according to claim 1, wherein:
each R is the same; and R3 = R4.

10

5. A bleaching composition according to claim 1, wherein
R3 and R4 are the same and are $-(CH_2)_n C(O)O-C1-C4-alkyl$.

15

6. A bleaching composition according to claim 1, wherein
R3 and R4 are selected from the group consisting of $-CH_2OH$,
 $-C(O)O-C1-C6-alkyl$, and phenyl.

20

7. A bleaching composition according to claim 1, wherein
at least one R1 and R2 is a 3-C0-C6-alkyl-pyridin-2-yl-C0-
C6-alkyl.

8. A bleaching composition according to claim 1, wherein Y
= 1

25

9. A bleaching composition according to claim 1, wherein
R3 and R4 are $-C(O)O-C1-C6-alkyl$.

30

10. A bleaching composition according to claim 1, wherein
at least one of R1 and R2 is selected from the group
consisting of: 3-ethyl-pyridin-2-ylmethyl, pyridin-2-

- 41 -

ylmethyl, 3-methyl-pyridin-2-ylmethyl, and 6-amide-pyridin-2-ylmethyl.

11. A bleaching composition according to claim 10, wherein
5 at least one of R1 and R2 is pyridin-2-ylmethyl.

12. A bleaching composition according to claim 1, wherein
both R1 and R2 are pyridin-2-ylmethyl and R is H.

10 13. A bleaching composition according to claim 1, wherein X
is C=O.

14. A bleaching composition according to claim 1, wherein
the bleaching composition comprises the free ligand.

15 15. A bleaching composition according to claim 1, wherein
the complex is of the general formula (A1):



20 in which:

M represents a metal selected from Mn(II)-(III)-(IV)-(V), Cu(I)-(II)-(III), Fe(II)-(III)-(IV)-(V), Co(I)-(II)-(III), Ti(II)-(III)-(IV), V(II)-(III)-(IV)-(V), Mo(II)-(III)-(IV)-(V)-(VI) and W(IV)-(V)-(VI);

X represents a coordinating species selected from any mono, bi or tri charged anions and any neutral molecules able to coordinate the metal in a mono, bi or tridentate manner;

30 Y represents any non-coordinated counter ion;
a represents an integer from 1 to 10;

- 42 -

k represents an integer from 1 to 10;

n represents an integer from 1 to 10;

m represents zero or an integer from 1 to 20; and

L represents a ligand as defined in claims 1 to 12, or
5 its protonated or deprotonated analogue.

16. A bleaching composition according to claim 15, wherein
M represents a metal selected from Fe(II) - (III) - (IV) - (V).

10 17. A bleaching composition according to claim 16, wherein
M represents a metal selected from Fe(II) and Fe(III).

18. A ligand of formula (I) according to claim 1 or a
transition metal catalyst thereof with the proviso that the
15 following compounds are excluded:

dimethyl 2,4-di-(2-pyridyl)-3,7-bis-(pyridin-2-ylmethyl)-
3,7-diaza-bicyclo[3.3.1]nonan-9-one-1,5-dicarboxylate;

1,5-bis-(hydroxymethylene)-2,4-di-(2-pyridyl)-3,7-bis-
(pyridin-2-ylmethyl)-3,7-diazabicyclo[3.3.1]nonan-9-ol;

20 dimethyl 2,4-di-(2-pyridyl)-3,7-bis-(pyridin-2-ylethyl)-3,7-
diaza-bicyclo[3.3.1]nonan-9-one-1,5-dicarboxylate;

dimethyl 2,4-di-(2-pyridyl)-3-(5-carboxypentyl)-7-methyl-
3,7-diaza-bicyclo[3.3.1]nonan-9-one-1,5-dicarboxylate;

25 dimethyl 2,4-di-(2-pyridyl)-3-(2-methoxyethyl)-7-methyl-3,7-
diaza-bicyclo[3.3.1]nonan-9-one-1,5-dicarboxylate; diethyl-

2,4-dipyridyl-7-picolyl-3,7-diaza-bicyclo-[3.3.1]-nonan-9-
one-1,5-dicarboxylate ; diethyl-2,4-dipyridyl-7-benzyl-3-

hydroxyethyl-3,7-diaza-bicyclo-[3.3.1]-nonan-9-one-1,5-
dicarboxylate; and, dimethyl-2,4-dipyridyl-7-benzyl-3-

30 hydroxyethyl-3,7-diaza-bicyclo-[3.3.1]-nonan-9-one-1,5-
dicarboxylate.

- 43 -

19. A ligand of formula (I) according to claim 18 or a transition metal catalyst thereof, wherein at least one of R1 or R2 is pyridin-2-ylmethyl and the other is selected from -CH3, -C2H5, -C3H7, and -C4H9.

5

20. A perchlorate salt of dimethyl 2,4-di-(2-pyridyl) -3,7-di(pyridin-2-ylmethyl) -3,7-diaza-bicyclo[3.3.1]nonan-9-one-1,5-dicarboxylate (N2Py4).

100434-434
100434-434